

Con. 7885-13.

GX-12071

(3 Hours)

[Total Marks : 80]

- N.B. : (1) Question no. 1 is **compulsory**.
 (2) Attempt any **three** questions out of the remaining **five** questions.
 (3) **Figures to right** indicate **Full marks**.

1. (a) Prove that real and imaginary parts of an analytic function $F(z) = u + iv$ are **5** harmonic function.
- (b) Find fourier series for $f(x) = |\sin x|$ in $(-\Pi, \Pi)$. **5**
- (c) Find the Laplace transform of $\int_0^t ue^{-3u} \sin 4u du$ **5**
- (d) If $\bar{F} = xy e^{2z} \hat{i} + xy^2 \cos z \hat{j} + x^2 \cos xy \hat{k}$, find $\operatorname{div} \bar{F}$ and $\operatorname{curl} \bar{F}$. **5**
2. (a) Using Laplace transform, solve :-
 $(\frac{D}{Dt}^2 + 3\frac{D}{Dt} + 2)y = e^{-2t} \sin t$ where $y(0) = 0, y'(0) = 0$. **6**
- (b) Find the directional derivative of $d = x^2 y \cos z$ at $(1, 2, \frac{\Pi}{2})$ in the direction of **6**
 $\hat{r} = 2\hat{i} + 3\hat{j} + 2\hat{k}$
- (c) Find the fouries series expansion for $F(x) = \sqrt{1 - \cos x}$ in $(0, 2\Pi)$, Hence deduce **8**
 that $\frac{1}{2} = \sum \frac{1}{4^n - 1}$.
3. (a) Prove the $J_{\frac{1}{2}}(x) = \sqrt{\frac{2}{\Pi x}} \left\{ \frac{\sin x}{x} - \cos x \right\}$. **6**
- (b) Evaluate by green's theorem, $\oint_C (x^2 y dx + y^3 dy)$ Where C is the closed path formed **6** by $y = x, y = x^2$.
- (c) (i) Find Laplace transform of $\frac{\cos bt - \cos at}{t}$ **4**

Con. 7885-GX-12071-13.**2**(ii) Find Laplace transform of :- $\frac{d}{dt} \left[\frac{\sin t}{t} \right]$

4

4. (a) Show that the set of functions $\{\sin x, \sin 3x, \dots\}$ OR $\{\sin(2n+1)x : n = 0, 1, 2, \dots\}$ is orthogonal over $[0, \frac{\pi}{2}]$, Hence construct 6 orthonormal set of functions.

(b) Find the imaginary part whose real part is $u = x^3 - 3xy^2 + 3x^2 - 3y^2 + 1$

(c) Find inverse Laplace transform of :-

6

(i) $\log \left(\frac{s^2 + 4}{s^2 + 9} \right)$

8

(ii) $\frac{s}{(s^2 + 4)(s^2 + 9)}$

5. (a) Obtain half range sine series for $f(x) = x^2$ in $0 < x < 3$.

6

- (b) A vector field \bar{F} is given by $\bar{F} = (x^2 - yz)\hat{i} + (y^2 - zx)\hat{j} + (z^2 - xy)\hat{k}$ is irrotational and Hence find scalar point function ϕ such that $\bar{F} = \nabla\phi$

6

- (c) Show that the function $V = e^x (x \sin y + y \cos y)$ satisfies Laplace equation and find its corresponding analytic function and its harmonic conjugate.

8

6. (a) By using stoke's theorem, evaluate $\oint_C [(x^2 + y^2)\hat{i} + (x^2 - y^2)\hat{j}] \cdot d\bar{r}$ where 'C' is the boundary of the region enclosed by circles $x^2 + y^2 = 4$, $x^2 + y^2 = 16$.

6

- (b) Show that under the transformation $w = \frac{5-4z}{4z-2}$ the circle $|z| = 1$ in the z-plane is transformed into a circle of unity in the w-plane.

6

- (c) Prove that $\int J_3(x) dx = \frac{-2J_1(x)}{x} - J_2(x)$.

8